

Tips for Living with Atrial Fibrillation

Atrial fibrillation is a type of irregular heartbeat that is often rapid. It is a common kind of *arrhythmia*—a condition in which your heart beats with an abnormal rhythm. However, many older people with atrial fibrillation don't have any obvious symptoms. Atrial fibrillation can be intermittent (occurs off-and-on) or chronic (present all the time).

Common Symptoms

- Palpitations
- Racing or fluttering of the heart
- Shortness of breath
- Chest discomfort
- Lightheadedness and dizziness
- Weakness or tiredness
- Anxiety

Common Causes

- High blood pressure
- Prior heart attack
- Heart valve problems
- Other heart disorders

The risk of atrial fibrillation increases with age, and it is present in about 10% of people 80 years of age or older. It is somewhat more common in men than in women. It is also more common in people with overactive thyroid, sleep apnea, emphysema, and other lung diseases. Atrial fibrillation is a leading cause of strokes in older adults.

Here's what the experts at the Health in Aging Foundation recommend:

GET REGULAR CHECK-UPS.

If you have atrial fibrillation, it is important to see your healthcare provider regularly. Discuss any new symptoms or if your treatment isn't controlling your symptoms. If so, the provider can change your treatment to help you feel better.

TAKE YOUR MEDICINES AS PRESCRIBED.

Medicines for atrial fibrillation work to keep the heart rate under control, reduce symptoms, and decrease the risk of stroke. In order to work properly and provide the greatest protection, medicines must be taken regularly and according to the instructions. Ask your healthcare provider if you have any questions.

TALK TO YOUR PHARMACIST OR HEALTHCARE PROVIDER.

Ask your pharmacist or healthcare provider **BEFORE** you take a new prescription, over-the-counter drug, herbal medicine, or other supplement. He or she can tell you about possible side effects and interactions with other medications you are already taking.

**LEARN THE
WARNING
SIGNS OF
STROKE.**

Because atrial fibrillation can increase your risk of having a stroke, it is very important that you recognize the warning signs of stroke:

- sudden numbness or weakness on one side of your face, arm, or leg
- slurred speech, garbled speech, or difficulty finding words
- sudden difficulty with balance or coordination
- sudden loss of vision, blurred vision, or double vision

Less common signs of stroke include:

- severe headache or sudden dizziness
- sudden confusion or change in level of awareness

Call 911 if you have any of these symptoms.

**IF YOU ARE
TAKING
WARFARIN,
BE CAREFUL
ABOUT EATING
CERTAIN
FOODS.**

This medication (also known by its brand name, Coumadin®) decreases the risk of stroke by making your blood less likely to form clots. (Warfarin is sometimes referred to as a “blood thinner”, but it does not actually thin the blood.)

Some foods can make warfarin stronger or weaker, especially certain vegetables and vegetable oils that are high in vitamin K. Examples are lettuce, cabbage, broccoli, and other greens. These vegetables are good for your health, so don't skip them. However, changing how much of these vegetables you eat can interfere with warfarin and increase the risk of stroke or bleeding. Instead, try to eat a similar amount on a regular basis and talk to your healthcare provider about *how* you can eat them safely.

**QUIT SMOKING
AND AVOID
EXCESS
ALCOHOL.**

The nicotine in tobacco can increase your risk of atrial fibrillation. Smoking also damages the heart and lungs and increases the risk of cancer. If you smoke, quitting is one of the best things you can do for your health. If you need help quitting, talk to your healthcare provider.

For some people, alcohol can increase the risk of atrial fibrillation, especially if you drink large amounts.

If alcohol triggers your atrial fibrillation, you should avoid it completely.

**ENGAGE IN
PHYSICAL
ACTIVITY.**

Regular activity such as walking and cycling can make your heart stronger and lower your risk of heart problems. It can also help you lose extra weight—a healthy weight is an important part of managing atrial fibrillation. Work with your healthcare provider to come up with an exercise plan that will be safe, effective, and enjoyable.