



COVID-19 Vaccines for Adults (age 65+)

Protect yourself against COVID-19. Be up to date with your vaccines!

You are **up to date with your COVID-19 vaccine** if you have gotten your primary series of COVID-19 shots and a bivalent booster (the updated booster).

The bivalent COVID-19 vaccines protect us against two different strains of COVID-19. The only COVID-19 booster vaccines approved by the FDA (as of September 8, 2022) are the Pfizer-BioNTech (age 12+) and Moderna (age 18+) bivalent vaccines.

FDA Approved Vaccine Shots for Older Adults (65+)

Vaccine	Primary Series	Bivalent Boosters
Pfizer-BioNTech	2 shots, 3 weeks apart	1 shot, at least 2 months after your last primary series shot or any other booster shots you received.
Moderna	2 shots, 4 weeks apart	
Novavax	2 shots, 3 weeks apart	Pfizer-BioNTech or Moderna
Johnson & Johnson/ Janssen*	1 shot of J&J/Janssen vaccine - 1 booster shot of either Moderna or Pfizer-BioNTech at least two months after first shot of J&J/Janssen	

* In most cases, Pfizer-BioNTech, Moderna, or Novavax vaccines are recommended over the Johnson & Johnson vaccine.

Approved COVID-19 Vaccines for Immunocompromised Adults (age 65 +)

A person is immunocompromised when their immune system is weak. These people may have more trouble fighting COVID-19 and are at higher risk of getting seriously sick or dying from COVID-19. You may have a weak immune system if you have cancer, diabetes, or heart conditions, for example. These are just a few of the diseases that can make your immune system weak. Please check with your healthcare provider to see if you have a medical problem that makes your immune system weak.

FDA Approved Vaccine Shots for Immunocompromised Older Adults (65+)

Vaccine	Primary Series	Bivalent Boosters
Pfizer-BioNTech	3 shots -2nd shot given 3 weeks after 1st shot -3rd shot given at least 4 weeks after 2nd shot	1 shot, at least 2 months after your last primary series shot or last booster Must be Pfizer-BioNTech
Moderna	3 shots -2nd shot given 4 weeks after 1st shot -3rd shot given at least 4 weeks after 2nd shot	1 shot, at least 2 months after your last primary series shot or last booster Pfizer-BioNTech or Moderna
Novavax	2 shots, 3 weeks apart	
Johnson & Johnson/ Janssen*	2 shots, 4 weeks apart - 2nd shot should be Pfizer or Moderna	

*In most cases, Pfizer, Moderna, or Novavax vaccines are recommended over the Johnson & Johnson vaccine.



DID YOU KNOW?

The vaccine is available at no cost to you:

- If you have Medicare, you will not have to pay to get vaccinated.
- Medicaid and private health insurance plans cover all vaccine costs.
- If you are uninsured you can get free vaccines.

Vaccines are available at many locations:

- Most drug stores
- Community health centers
- State or local health departments

Find a COVID-19 vaccine location near you:

- Ask your doctor or healthcare provider
- Call 1-800-232-0233
- Text your zip code to 438829 or visit [vaccines.gov](https://www.vaccines.gov)



40 FULTON STREET
SUITE 809
NEW YORK, NY 10038
212.308.1414 TEL
Info@healthinaging.org

The AGS Health in Aging Foundation is dedicated to improving the health, independence, and quality of life of all older people. We aim to empower older adults and caregivers to actively participate in their health care and decision-making.

DISCLAIMER: This information is not intended to diagnose health problems or to take the place of medical advice or care you receive from your physician or other healthcare provider. Always consult your healthcare provider about your medications, symptoms, and health problems. September 2022

©2022 Health in Aging Foundation. All rights reserved.