# HealthinAging.org

Trusted Information. Better Care.

Expert Information from Healthcare Professionals Who Specialize in the Care of Older Adults



## **Essential Vaccination Information for Older Adults**

Some factors that affect your health are outside of your control. However, many important risk factors are within your power to change. This includes getting shots, called vaccinations, that help protect you from certain illnesses. Vaccines are some of the safest therapies around. Although all therapies, including vaccines, pose the rare chance of serious side effects, for most people, the risks from the diseases are far greater than the risks from the vaccines. Contact your health care provider or local health department for a list of doctors who give these shots. Your city or county health department or local hospital may operate clinics that provide these vaccines. The American Geriatrics Society's Health in Aging Foundation recommends the following vaccinations for most older adults.

#### **Flu Shot**

What it does	Protects against annual influenza viruses.
Who needs it	The Centers for Disease Control and Prevention (CDC) recommends any type of flu shot licensed for persons aged ≥65 years (standard or high-dose, adjuvanted or unadjuvanted or recombinant influenza vaccines). While everyone should get a flu vaccine, the CDC notes that it is especially important for the following people to get flu shots because they are at high risk for having serious flu-related complications: anyone who is 65 years of age or older; nursing home residents; and people with serious health conditions such as heart disease, diabetes, asthma, lung disease or HIV. Caregivers for older adults should also get vaccinated to avoid spreading the flu.
Who should not get it	People who have had allergic reactions to flu shots in the past, or have been diagnosed with Guillian-Barre Syndrome.
When to get it	Because new strains of the flu develop constantly, the flu vaccine must be given yearly. You should get your flu shot in the fall.

#### **Pneumococcal Shot**

What it does	Protects against pneumococcal bacteria, which can cause pneumonia and blood and brain infections.
Who needs it	Anyone 65 years or older.
When to get it	Ask your healthcare provider. Two different types of pneumococcal vaccine are recommended now. They are called pneumococcal conjugate vaccine (PCV)13 and pneumococcal polysaccharide vaccine (PPSV)23.

### **Tetanus/Diphtheria Shot**

What It Does	Protects against two potentially deadly bacterial infections. A second, and different, form of the vaccine (called "Tdap") protects against tetanus, diphtheria, and pertussis (whooping cough).
Who Needs It	Everyone. It is now recommended to get a one-time dose of the Tdap version if you are 65 or older to be protected from whooping cough.
When to get it	Once every 10 years.

#### **Shingles (Herpes Zoster) Shot**

#### What It Does

**t Does** A new shingles (herpes zoster) vaccine, also known as Recombinant Zoster Vaccine (RZV) and the older live zoster vaccine protect against the development of shingles. The recombinant zoster vaccine reduces the risk by 97% in persons 50 years of age and older, and about 90% in persons 70 years of age and older. The RZV protects against the development of chronic pain from shingles (also called postherpetic neuralgia), reducing the risk by 85% in persons 70 years of age and older. The live zoster vaccine reduces the risk of shingles by 70% in persons age 50-59, 64% age 60-69, 37% age 70 and older, and reduces postherpetic neuralgia by 66% across all age groups.

**Who Needs It** The CDC recommends the recombinant zoster vaccine over the live zoster vaccine for persons age 50 years and older.

Who ShouldPeople who have a history of a severe allergic reaction to any component of the vaccine.Not Get It

## When to get itAsk your healthcare provider. The CDC also recommends the recombinant zoster vaccinefor persons who previously received the live vaccine.

The recombinant zoster vaccine requires two doses with the second dose given anytime between 2–6 months after the first dose.

**The Centers for Disease Control and Prevention (CDC) also recommends additional shots** for older adults who run an increased risk of these diseases because they have certain health problems, occupations, or lifestyles. These include the Measles, Mumps, Rubella (MMR) vaccination, and shots for Varicella, Hepatitis A and B, and Meningococcal disease. Ask your healthcare provider if you should get any of these additional shots. For additional information, visit the CDC website at www.cdc.gov.

HEALTH IN AGING F O U N D A T I O N THE OFFICIAL FOUNDATION OF THE AMERICAN GERIATRICS SOCIETY 40 FULTON STREET 18TH FLOOR NEW YORK, NY 10038 212.308.1414 TEL 212.832.8646 FAX Info@healthinaging.org *DISCLAIMER*: This information is not intended to diagnose health problems or to take the place of medical advice or care you receive from your physician or other healthcare provider. Always consult your healthcare provider about your medications, symptoms, and health problems. Sept 2019.

©2019 Health in Aging Foundation. All rights reserved. This material may not be reproduced, displayed, modified, or distributed without the express prior written permission of the copyright holder. For permission, contact info@healthinaging.org.